

Astronomical Site Ranking Based on Tropospheric Wind Statistics

B. García-Lorenzo^{1*}, J.J. Fuensalida¹, C. Muñoz-Tuñón¹, and E. Mendizabal^{1,2}

¹ Instituto de Astrofísica de Canarias, C/Via Lactea S/N, 38350-La Laguna, Tenerife, Spain

² Instituto Nacional de Astrofísica, Óptica y Electrónica, C/Luis Enrique Erro,1, Tonantzintla, Puebla 72840, Mexico

Accepted Received; in original form

ABSTRACT

High altitude wind speeds have been adopted as a parameter for astronomical site selection based on the relationship found at the Paranal and Cerro Pachón sites between the average velocity of the turbulence (V_0) and winds at the 200 millibar pressure level (V_{200}). Although this relationship has not been checked at any other site in the world and a connection between image quality and V_{200} has not been proven anywhere, high altitude wind speed (V_{200}) is a parameter for checking the suitability of sites for adaptive optics and surveying potential sites for ELT.

We present comprehensive and reliable statistics of high altitude wind speeds and the tropospheric flows at the location of five important astronomical observatories. We have used the National Center for Environmental Prediction (NCEP/NCAR) Reanalysis database to collect wind data at nine tropospheric pressure levels for the five selected sites. For comparison and validation of the data from the climate diagnostic model, we have also obtained wind profiles from radiosonde stations. The degrees of correlation found indicate a high level of significance between NCEP/NCAR Reanalysis and balloon databases, pointing to NCEP/NCAR Reanalysis as a useful database for site characterization.

Statistical analysis exclusively of high altitude winds point to La Palma as the most suitable site for adaptive optics, with a mean value of 22.13 m s^{-1} at the 200 mbar pressure level. La Silla is at the bottom of the ranking, with

the largest average value 200 mbar wind speed (33.35 m s^{-1}). We have found a clear annual periodicity of high altitude winds for the five sites in study.

We have also explored the connection of high to low altitude atmospheric winds as a first approach of the linear relationship between the average velocity of the turbulence and high altitude winds (Sarazin & Tokovinin 2001). We may conclude that high and low altitude winds show good linear relationships at the five selected sites. The highest correlation coefficients correspond to Paranal and San Pedro Mártir, while La Palma and La Silla show similar high to low altitude wind connection. Mauna Kea shows the smallest degree of correlation, which suggests a weaker linear relationship. Our results support the idea of high altitude winds as a parameter for rank astronomical sites in terms of their suitability for adaptive optics, although we have no evidence for adopting the same linear coefficient at different sites. The final value of this linear coefficient at a particular site could drastically change the interpretation of high altitude wind speeds as a direct parameter for site characterization.

Key words: Site Testing — Turbulence — Instrumentation: Adaptive Optics

1 INTRODUCTION

Ground-based stronomical observations are drastically affected by the presence of atmospheric turbulence. The theoretical angular resolution of a telescope, determined by its diameter, is usually severely degraded because of atmospheric turbulence. Adaptive optics techniques have been developed to compensate for the effects of the atmosphere on astronomical images and to reach the diffraction limit of our telescopes. However, the design and operation of adaptive optic systems are complex and difficult because the turbulence above astronomical sites is still insufficiently well characterized. In spite of partial information on the turbulence behaviour above existing astronomical observatories, it is obvious that a site with low and stable turbulence should be better suited for the implementation of adaptive optics than one with a turbulent and chaotic atmosphere. The requirements for excellent image quality of current large and future very large telescopes demand a proper knowledge of atmospheric turbulence, and several projects are already pursuing this aim. Moreover, the precise characterization of the turbulence above a particular site requires long term monitoring. To counter the lack of long term information on turbulence, high altitude winds (in

particular winds at the 200 mbar pressure level—hereafter V_{200}) have been proposed (Sarazin & Tokovinin 2001) as a parameter for estimating the total turbulence at a particular site, since records of this parameter exist from several sources. This proposal is based on the idea that the greatest source for turbulence generation is related to the highest peak in the vertical wind profile, which is located at the 200 mbar pressure level globally. Also, Sarazin & Tokovinin (2001) find a good correlation between the average velocity of the turbulence, V_0 , and V_{200} of the form $V_0 = 0.4V_{200}$ at the Cerro Pachón and Paranal Observatories in Chile. Although the $V_0 \propto V_{200}$ relationship has not been tested at other sites, its validity would simplify the calculation of key parameters for adaptive optics, such as the coherence time. Furthermore, V_{200} statistics could be used as a parameter for ranking the suitability of different observing sites for AO. However, such a relationship never indicates a connection between the total turbulence at the ground level (seeing) and wind speed at the 200 mbar pressure level. Indeed, Sarazin & Tokovinin (2001) have also studied this option but a connection has not been found. Despite this, a false idea of a connection between image quality and high altitude wind speed is increasingly widespread among those in the astronomical community interested in adaptive optics.

Some studies centred on V_{200} statistics values have been recently published (Carrasco & Sarazin 2003; Sarazin 2002), and a detailed statistical analysis of V_{200} for Roque de los Muchachos Observatory (ORM) on the island of La Palma (Spain) have been performed by Chueca et al. (2004). While the linear $V_0 \propto V_{200}$ relationship (Sarazin & Tokovinin 2001) is confirmed at other sites, a first approach to connecting high altitude winds to turbulence at ground level can be carried out by studying the wind vertical profile and the relation of high to low altitude winds. Previous studies have also reported a connection of ground layer winds to image quality (Muñoz-Tuñón, Varela & Mahoney (1998); Varela, Muñoz-Tuñón, & Gurtubai (2001)).

In this paper we present a detailed statistical analysis of V_{200} for different observing sites in the two hemispheres. For the first time, we study the connection of high to low altitude winds and we present the statistical analysis of the wind vertical profile at five astronomical sites, the ORM, Mauna Kea and San Pedro Martir in the north, and Paranal and La Silla in the south (Table 1). We also discuss the suitability for adaptive optics of the five summits analysed on the basis of the wind vertical profile statistics.

Table 1. Location of selected observing sites (from <http://www.seds.org/billa/bigeyes.html>)

	ORM(La Palma)	La Silla	Mauna Kea	Paranal	San Pedro Mártir
Latitude	28 46 N	29 15 S	19 50 N	24 38 S	31 02 N
Longitude	17 53 W	70 44 W	155 28 W	70 24 W	115 27 W
Altitude	2400 m	2400 m	4100 m	2635 m	2800 m
Country	Canary Islands, Spain	Chile	Hawaii, USA	Chile	Baja California, Mexico

Table 2. Radiosonde stations closest to the selected astronomical sites.

Station	Place	Latitude	Longitude	Available Range	Close to Site	Distance to Site (km)
600200	Santa Cruz de Tenerife (Spain)	28.467 N	16.25 W	1960-2001	La Palma	163
855430	Quintero (Chile)	32.780 S	71.517 W	1957-1999	La Silla	400
912850	Hilo (Hawaii, USA)	19.716 N	155.06 W	1950-2002	Mauna Kea	44.5
854420	Antofagasta (Chile)	23.410 S	70.467 W	1957-2001	Paranal	136
722930	California (USA)	32.867 N	117.15 W	1972-2001	San Pedro Martir	260

2 THE DATA

We have selected five existing sites spread all over the globe to study the behaviour of V_{200} and the vertical wind profile. Table 1 lists the astronomical observatories considered, their location and altitude. We used the National Center for Environmental Prediction (NCEP/NCAR) Reanalysis database to collect wind speeds at different pressure levels for the five sites. The Reanalysis data were obtained as 6-hourly and daily U -wind and V -wind components from the beginning of 1980 to the end of 2002. Wind speeds in this database are considered as one of the most reliably analysed fields (see Kalnay et al. (1996) and Kistler et al. (2001) for a detailed description) and they are heavily constrained by observational data. eW obtained the 6-hourly and daily wind speed module from U -wind and V -wind 6-hourly and daily components (more than 33 600 data for each component and location).

For comparison and validation of the data from the climate diagnostic model, we have obtained winds from radiosonde stations. The main criteria in selecting radiosonde stations were the year range for available data and the distance from the observing sites. Table 2 shows information about the final selected stations.

In both databases we have selected the simultaneous data for the period 1980–2002 to derive the mean monthly wind at the different common pressure levels. We found a good correlation between the balloon and NCEP/NCAR data statistic in the sites studied. Figure 1 shows the linear Pearson correlation coefficient of the mean monthly V_{200} time series from the two databases at different pressure levels. The degrees of correlation found indicate a high level of significance between the NCEP/NCAR Reanalysis and balloon databases at




Figure 1. Linear Pearson correlation coefficient obtained comparing the mean monthly wind statistics for the NCEP/NCAR data and balloon measurements at different pressure levels for the period 1980–2002. The height axes have been derived from the pressure to height relationship assuming ICAO standard atmosphere.

the five selected locations and for any altitude considered, thereby revealing NCEP/NCAR Reanalysis as a useful database for site characterization approach. The detailed analysis of winds in the following sections will be performed only with the NCEP/NCAR Reanalysis data at each site. This database provide a better temporal coverage and resolution than the balloon measurements and have no gaps in the time series

3 ANALYSIS

Sarazin & Tokovinin (2001) have suggested that the suitability of a site for adaptive optics is related to winds at the pressure level of 200 mbar. The average altitude of this pressure level at the Earth’s poles is about one kilometre lower than at the equator. Moreover, the 200 mbar pressure level presents a slightly seasonal altitude dependency, being about 400 m lower in altitude in winter than in summer time. All other pressure levels show a similar behaviour. Assuming an ICAO standard atmosphere (see <http://www.pdas.com/coesa.htm>

for an exact definition), we shall consider hereafter a mean altitude of 12170 meters for the 200 mbar pressure level at any site. In the same way, we adopt the same mean altitude for any of the pressure levels considered in this paper at any site. Therefore, the height axis of the figures can only be taken as an approximate value.

3.1 V_{200} statistics and seasonal behaviour

We have derived the monthly average, V_{200} , for the period 1980–2002 for the five selected sites, and the resulting monthly statistics are shown in figure 2A. There is a seasonal trend in the high altitude wind behaviour. In the northern hemisphere, the highest V_{200} occurs during spring and the lowest in summer. La Silla and Paranal in the southern hemisphere show high V_{200} values during the southern winter and spring.

In the northern hemisphere, San Pedro Mártir shows the lowest V_{200} during the summer, while the lowest V_{200} in winter corresponds to La Palma. In autumn, Mauna Kea and La Palma show similar values of V_{200} . The differences in V_{200} are small for the three northern observatories during the spring. We define V_N as the ratio of the monthly average wind velocity at 200 mbar to the V_{200} mean over the full period (Table 3) ($V_N = V_{200\text{monthly}}/V_{200\text{globalmean}}$). This ratio has the same seasonal behaviour as V_{200} at each observatory but magnitudes are related to the same reference level, the V_N mean, which is always equal to unity. If S_{200} is the monthly standard deviation of V_{200} , the V_N to S_{200} ratio gives a view of the stability of high altitude winds, as shown in Figure 2B. The V_N/S_{200} ratio at any site are directly comparable, with higher V_N/S_{200} values indicating more stable V_{200} regimes. According to the defined stability parameter, V_N/S_{200} , San Pedro Mártir presents the least stable high altitude winds of the three northern sites. La Palma shows a similar stability parameter throughout the year, with the more stable V_{200} during May and June. Mauna Kea presents a seasonal behaviour of the stability parameter, with highest values in February and March and lowest during May and June. Although the high altitude winds at Mauna Kea are more stable than at La Palma from January to March, V_{200} is higher in Mauna Kea during these months. Only September and October shows a better behaviour at Mauna Kea than at La Palma, with similar V_{200} values but high stability at Mauna Kea.

In the south hemisphere, Paranal has more stable high altitude winds than La Silla except in January. At Paranal, November is the most stable month in terms of V_{200} . Table 3

Table 3. Statistical results of 200 mbar wind speed (m s^{-1}) by using the data from NCEP/NCAR Reanalysis database for the period 1980–2002 at different astronomical sites. Amplitude is the difference of maximum and minimum statistical winds.

Site	Amplitude (m s^{-1})	Mean (m s^{-1})	Median (m s^{-1})	Standard deviation (m s^{-1})	RMS (m s^{-1})
La Palma	13.69	22.13	20.79	11.67	0.06
La Silla	12.46	33.35	32.77	12.94	0.07
Mauna Kea	18.00	24.33	22.81	12.30	0.07
Paranal	18.47	30.05	28.63	13.01	0.07
San Pedro	26.49	26.55	24.57	15.39	0.08




Figure 2. (A) The monthly average wind velocity at 200-mbar pressure level for the period 1980–2002 for the five selected observing sites. (B) Normalized mean to standard deviation ratio of wind velocity at 200-mbar pressure level for the period 1980–2002 for the selected observing sites. In panels, lines and symbols indicate sites as in figure 1.

lists the V_{200} mean, median, standard deviation, and root mean square values for the period 1980–2002 for the five observatories studied.

In terms of V_{200} , La Palma is the best site of the five studied, with a mean value of 22.13 m s^{-1} , the smallest standard deviation (Table 3) and showing the most stable high altitude winds. In contrast, La Silla shows the highest average value, with a mean V_{200} of 33.35 m s^{-1} , although the amplitude in Table 3 (the difference between minimum and maximum) at this site is only 12.46 m s^{-1} , the lowest of the five observatories studied. The statistical V_{200} amplitudes for Paranal and Mauna Kea are quite similar, while the value for San Pedro Mártir is approximately twice the V_{200} amplitude at La Silla or La Palma. According to the relationship connecting V_0 and V_{200} (Sarazin & Tokovinin 2001), large V_{200} amplitudes indicate widely varying turbulence conditions over the course of the year.

In order to determine any dominant temporal variation of V_{200} over the observatories in question, we have performed a wavelet analysis (see Torrence & Compo (1998) for a practical

step-by-step description of wavelet analysis) of the time series obtained from NCEP/NCAR databases for the five selected observatories. We used a Morlet function as wavelet mother. Figure 3 shows the time series (plot a), the local wavelet power spectrum (plot b) and the time average power spectrum (plot c) corresponding to the wavelet analysis of each of the five observatories in study. A table of the reconstruction time series accuracy for each site is shown in panel F of Figure 3. The wavelet analysis of V_{200} for La Palma site has already been discussed by Chueca et al. (2004), and is in good agreement with the present results.

The annual periodicity of V_{200} remains significant for all the five sites studied. Except for Mauna Kea, a 0.5 years period appears in the global power spectrum for the other four selected sites, although only for La Palma is this peak still significant in the local wavelet power. For La Palma and La Silla, a 2.3 years period is also significant in the local power spectrum, but they are less than significant globally. A four year period appears for the five observatories in study, but only in San Pedro Mártir does it remain significant in the global power spectrum.

Atmospheric conditions at the Paranal site has changed considerably from site testing period in the early 1990s to the present (ESO, 2001). The increase in the average seeing from 1998 onwards at Paranal has been identified with El Niño/La Niña events in South America (Beniston, Casals, & Sarazin 2002). However, we have found no similar discrepancy between high altitude winds before and after 1998 at Paranal. The temporal evolution of the annual periodicity of V_{200} remains quasi-constant from 1990 (Figure 3D). The mean and standard deviation of V_{200} for the period 1990–4 are 29.19 m s^{-1} and 7.66 m s^{-1} , respectively. These values are slightly lower than those obtained for the period 1998–2002 (30.77 m s^{-1} and 8.31 m s^{-1} for the mean and standard deviation of V_{200} , respectively) but they are still in agreement. Therefore, the influence of events affecting the atmospheric conditions at Paranal site at low altitude has no counterpart to winds at high altitude.

3.2 High and low altitude wind connection

Although only wind speed at the 200 mbar pressure level has been proposed as a parameter for site characterization, it seems reasonable to assume that statistical wind profiles could also be important in determining stability and the connection of high altitude winds to turbulence at ground level. As a first approach, we explore the connection of high to lower altitude atmospheric winds by using the NCEP/NCAR database. We consider the pressure




Figure 3. Wavelet analysis results for the mean monthly 200 mbar wind velocity at: (A) ORM (La Palma), Spain; (B) La Silla, Chile; (C) Mauna Kea, Hawaii, USA; (D) Paranal, Chile; and (E) San Pedro Martir, Mexico. In these plots a) is the time series used for the wavelet analysis; b) is the local wavelet power spectrum of the time-series obtained from the Morlet wavelet. The colour scale at the bottom indicates the power spectrum intensity of the full range; plot c) is the average power spectrum. The dashed line is the 95 per cent confidence level. Panel (F) show the results comparing the reconstruction time series from the wavelet analysis and the input time series.

levels in the troposphere over the observing sites available from the NCEP/NCAR database. These pressure levels are 700, 600, 500, 400, 300, 250, 200, 150 and 100 millibars for La Palma, La Silla, Paranal and San Pedro Mártir. For Mauna Kea, the 700 mbar pressure level is always below the altitude of the observatory. The connection between the boundary layer and high altitude winds is the subject of a forthcoming study (Varela et al., in preparation).

In order to reveal the relationship of tropospheric winds at different altitudes, we have compared the 6-hourly and daily wind measurements at each pressure level with simultaneous data at any other levels. Figure 4 shows the results of this comparison. The lowest levels, 700 mbar (600 mbar for Mauna Kea), are most probably affected by the topography of the sites, which may be breaking the linear relationship. At Mauna Kea, the linear relationship of 150 and 200 mbar to 500 mbar is lower than 0.5, suggesting a weak connection between high and low altitude winds. At the other sites, there is a clear connection of high to low altitude winds, with Pearson coefficients greater than 0.5 for any level. These coefficients indicate a good linear relationship between the wind measurements at different pressure levels. Only the comparison of highest, 100 mbar, to the lowest, 700 mbar (600 mbar for Mauna Kea) level gives Pearson coefficients less than 0.5. Winds at the lowest levels could be affected by the surrounding site topography, perhaps disturbing the linear relationship. In fact, the 700 mbar pressure level corresponds to approximately 3120 metres, less than one kilometre above La Palma, La Silla, Paranal and San Pedro Mártir. For Mauna Kea, the mean altitude of the lowest pressure level, 600 mbar, is around 4350 metres, just 250 metres above the summit. At the other pressure extreme, 100 mbars corresponds on average to 16 750 metres, approximately at the average altitude of the tropopause layer (Seidel et al. 2001; García-Lorenzo, Fuensalida, & Eff-Darwich 2004). Therefore, the breaking of the linear relationship of 100 mbar pressure level to lower altitudes could be related to the fact that 100 mbar could be below the tropopause. In other words, winds measurements at 100 mbars could correspond to tropospheric, tropopausal or mesospheric winds, depending on the season.

Table 4 shows the mean values of winds at different levels, derived by averaging the individual Pearson coefficients obtained from comparing a particular level to any other (figure 4) and the standard deviation of these coefficients (σ_m). We have not considered the comparison of any level to itself (Pearson coefficient of 1 in Figure 4). Table 4 also includes the mean covariance (Cov.) and its standard deviation (σ_{cov}) as an indicator of the strongest of the linear relationships. Although the largest mean correlation coefficients corresponds to




Figure 4. Linear Pearson correlation coefficient obtained comparing the wind data at each pressure level to the wind measurements at any other level at: (a) La Palma, (b) La Silla, (c) Mauna Kea, (d) Paranal and (e) San Pedro Martir. The comparison level is indicated by a different line type or symbol according to the bottom-right table. The value equal to 1 also indicates the comparison pressure level.

Table 4. Mean linear relationship of wind measurements at different pressure levels. Mean corresponds to the average of Pearson coefficients obtained comparing the referred pressure level to any other. σ_m indicates the standard deviation of these coefficients. Cov. is the average covariance of individual covariance values obtained comparing each level with the others, while σ_{cov} is the standard deviation of these covariances.

Pressure (mbar)	La Palma				La Silla				Mauna Kea				Paranal				San Pedro			
	Mean	σ_m	Cov.	σ_{cov}	Mean	σ_m	Cov.	σ_{cov}	Mean	σ_m	Cov.	σ_{cov}	Mean	σ_m	Cov.	σ_{cov}	Mean	σ_m	Cov.	σ_{cov}
100	0.59	0.16	42.2	21.0	0.57	0.13	45.4	21.0	0.62	0.17	49.7	27.4	0.64	0.13	52.9	27.0	0.70	0.11	77.7	36.3
150	0.70	0.18	64.4	33.0	0.73	0.15	79.5	39.5	0.64	0.26	69.6	45.0	0.78	0.14	86.2	44.9	0.77	0.14	115.	57.8
200	0.74	0.18	76.2	38.0	0.78	0.15	97.7	47.4	0.67	0.27	76.3	48.4	0.78	0.17	98.1	49.7	0.79	0.14	132.	65.3
250	0.77	0.17	76.8	36.9	0.80	0.15	100.	47.5	0.71	0.23	74.3	43.4	0.83	0.14	97.0	48.3	0.81	0.13	131.	63.1
300	0.78	0.15	71.7	32.7	0.81	0.15	95.3	44.1	0.72	0.18	65.3	34.0	0.84	0.13	89.7	43.9	0.83	0.11	121.	55.6
400	0.79	0.12	58.2	22.7	0.81	0.14	78.7	33.6	0.68	0.13	43.6	16.1	0.83	0.12	72.3	32.7	0.83	0.09	97.8	38.8
500	0.76	0.14	45.5	14.4	0.78	0.14	61.3	23.1	0.58	0.19	26.6	5.80	0.81	0.12	55.8	22.2	0.81	0.11	76.4	25.6
600	0.68	0.18	33.1	8.34	0.74	0.16	45.4	14.3	0.40	0.22	13.8	3.10	0.76	0.13	38.2	12.4	0.76	0.13	54.8	14.5
700	0.55	0.20	22.1	4.84	0.60	0.17	28.1	7.32					0.63	0.15	20.1	4.70	0.67	0.15	34.9	6.55

400 mbar above La Palma, La Silla and San Pedro Mártir, and 300 mbar for Mauna Kea and Paranal, the maximum mean covariance corresponds to a level of around 200 mbar.

Combining Figure 4 and results of table 4, we may conclude that high altitude winds show good relationships to low altitude winds. The highest correlation corresponds to Paranal and San Pedro Mártir. La Palma and La Silla show a similar behaviour in this regard, while Mauna Kea shows the worse correlation between high to low winds. On the basis of these results, we deduce that the relationship $V_0 \propto V_{200}$ (Sarazin & Tokovin 2001) should be worse for Mauna Kea than for any of the other sites considered in this paper. The topographical influence seems to be slightly greater at the sites on islands than at the continental sites.

3.3 Wind profile statistics and seasonal behaviour

Figure 5 shows the monthly average wind for the period 1980–2002 at different pressure levels for the five observatories. Wind statistics at the different pressure levels follow a similar seasonal behaviour to that of V_{200} (Section 3.1), V_{200} being the highest winds at any month and site. Only Mauna Kea shows slightly stronger mean winds at 150 mbar than V_{200} during May. The monthly average winds at 100 mbar show the largest differences from the statistics of all the others at the pressure levels considered, perhaps because the 100 mbar level can be above or below the tropopause. The observatories studied seem to be statistically stable in the sense that they show the worst and best wind measurements for the same epoch at any of the pressure levels considered. Only Mauna Kea shows the strongest winds in levels at different months: while the statistical V_{200} peak corresponds to March, the strongest winds for 150 and 250 mbar occur in April and February, respectively. The monthly average wind behaviour shown in Figure 5 for Mauna Kea confirms the results




Figure 5. The monthly average wind velocity for the period 1980–2002 at the pressure levels indicated in the bottom-right table for the selected sites.

Table 5. Statistical results of wind speed profile (m s^{-1}) by using the data from NCEP/NCAR Re-analysis database for the period 1980–2002 at different astronomical sites. Amplitude is the difference of maximum and minimum statistical winds at each pressure level.

Site	Level (mbar)	Amplitude (m s^{-1})	Mean (m s^{-1})	Median (m s^{-1})	Standard deviation (m s^{-1})	RMS (ms^{-1})
La Palma	100	14.55	13.69	12.69	7.72	0.04
	150	14.39	20.51	19.43	10.24	0.06
	250	12.61	20.66	19.04	11.49	0.06
	300	11.30	18.23	16.56	10.49	0.06
	400	8.53	14.18	12.71	8.44	0.05
	500	5.29	11.64	10.42	6.86	0.04
	600	3.09	9.78	8.79	5.63	0.03
	700	3.02	8.27	7.52	4.64	0.03
La Silla	100	11.17	20.58	20.37	7.73	0.04
	150	10.68	31.09	30.74	10.93	0.06
	250	12.42	31.04	30.23	13.05	0.07
	300	12.06	27.52	26.70	12.20	0.07
	400	10.92	20.45	19.48	10.01	0.05
	500	9.32	14.80	13.73	7.96	0.04
	600	7.50	10.46	9.35	6.27	0.03
	700	5.18	7.20	6.16	4.68	0.03
Mauna Kea	100	14.14	12.76	10.85	8.35	0.05
	150	16.81	22.22	20.54	11.53	0.06
	250	16.98	21.39	19.77	11.35	0.06
	300	15.83	17.41	15.73	9.83	0.05
	400	11.88	11.27	9.75	7.12	0.04
	500	8.09	8.00	6.85	5.25	0.03
	600	4.93	6.24	5.46	3.99	0.02
Paranal	100	15.45	18.05	17.83	8.35	0.05
	150	18.09	28.58	27.99	11.73	0.06
	250	17.29	27.38	25.76	12.66	0.07
	300	15.46	23.85	22.29	11.54	0.06
	400	12.41	17.13	15.70	9.29	0.05
	500	10.34	12.12	10.66	7.30	0.04
	600	7.67	8.39	7.20	5.31	0.03
	700	3.98	5.43	4.80	3.31	0.02
San Pedro	100	19.49	15.45	14.27	9.59	0.05
	150	26.01	23.94	22.77	13.54	0.07
	250	24.07	24.62	22.24	14.87	0.08
	300	21.15	21.44	18.88	13.45	0.07
	400	16.22	16.22	13.88	10.77	0.06
	500	12.32	12.83	10.82	8.63	0.05
	600	8.45	10.26	8.71	6.52	0.04
	700	5.40	7.86	6.93	4.67	0.03

of the previous section, showing that the connection between high and low altitude winds is not clear for the Mauna Kea site.

Table 5 shows the annual statistical results of the wind profiles for the five observatories. Vertical wind gradients are larger in winter than in summer, indicating stabler and hence less turbulent atmospheres in summer all the observatories studied.

4 DISCUSSION AND CONCLUSION

The connection between V_{200} and the average velocity of the turbulence, V_0 , found by Sarazin & Tokovinin (2001) at Paranal and Cerro Pachón has been accepted as a parameter for tackling astronomical site suitability for adaptive optics. The linear relationship between V_0 and the vertical wind profile, $V(h)$, is established by the definition of V_0 :

$$V_0 = \frac{\int_0^\infty C_N^2(h)V(h)dh}{\int_0^\infty C_N^2(h)dh}, \quad (1)$$

where $C_N^2(h)$ is the turbulence vertical profile. The high level of agreement between winds at any altitude demonstrated at Section 3.2 therefore suggests a relationship between V_0 and V_{200} at any of the observing sites studied. However, we have only demonstrated a connection of high and low altitude winds, but not a connection of high altitude winds and the average velocity of the turbulence. Even if such a connection were valid world-wide, it could be either linear or non-linear. In the mean time we consider a linear relationship at the sites studied similar to that found at Paranal and Cerro Pachón. Our results support the idea of considering V_{200} as a parameter for ranking astronomical sites with regard to their suitability for adaptive optics in spite of the lack of information on $C_N^2(h)$. However, the importance of deriving atmospheric turbulence profiles is still essential. $C_N^2(h)$ determines the coefficient of the linear fit connecting V_0 and V_{200} (A hereafter, $V_0 = A \times V_{200}$). For Paranal and Cerro Pachón this coefficient is $A = 0.4$ (Sarazin & Tokovinin 2001), but there are still no measurements for any other site. According to the results of this paper, we do not have any evidence for adopting the same coefficient at the different sites. Even in the hypothetical case of similar $C_N^2(h)$ at different summits (which is less than probable), the connection between high to low altitude winds reveals several differences at different sites, suggesting a different A coefficient in the linear fit. The final value of the linear coefficient at a particular site could drastically change the interpretation of V_{200} as a direct parameter for site characterization. If we consider a site with V_{200} statistics similar to Paranal and A adopts larger values than 0.4 at that particular site, we obtain larger V_0 values than for Paranal. In contrast, a smaller A value will indicate less atmospheric turbulence at that hypothetical site than at Paranal.

Another important point is the seasonal behaviour of V_{200} . Changes in V_{200} regimes in different seasons could have an important influence on the value of A , taking into account that we have presented here only the statistics of wind modules, and that we have not considered the wind direction, which could point to changing wind regimes with season. We could obtain different A values for different months; that is, A could also present seasonal

behaviour. To have the best parameterization of the average wind turbulence, V_0 , in term of V_{200} seems to be mandatory in any study of the seasonal behaviour of the linear coefficient A . Unfortunately, our information about A is very poor, with measurements only at Paranal and Cerro Pachón (Sarazin & Tokovinin 2001). If A were a constant coefficient and had a similar value to that found for Paranal and Cerro Pachón, V_0 would present a similar seasonal behaviour to that of V_{200} . In this hypothetical case, the statistical results obtained here for V_{200} would indicate that La Palma is at the top of the suitability ranking list for adaptive optics, while La Silla is at the bottom.

Atmospheric stability, in the sense of similar behaviour and good correlation of winds at different altitudes, is also an important point in establishing V_{200} as a direct parameter for site characterization and adaptive optics suitability. Our results show that Paranal and San Pedro Mártir present statistically more stable atmospheres than the other three sites considered. The stability results are found similar for La Silla and La Palma. Mauna Kea shows the worst results with regard to high to low altitude winds. Taking into account the behaviour of V_{200} and its connection to low altitude winds, Mauna Kea should present unstable average wind turbulence compared to the other sites. However, the stability of winds at different altitudes is not a guarantee of stable turbulence behaviour. If $C_N^2(h)$ is stable enough at a site with unstable winds behaviour in altitude, it should be a similar or better site for adaptive optics than a site with stable winds and unstable $C_N^2(h)$ behaviour. A more stable atmosphere, in the sense established in this paper, suggest a better relationship between V_0 and V_{200} and better accuracy of the determination of the A coefficient than sites with unstable atmospheres.

Although a correlation between seeing and winds at the 200 mbar pressure level has not been found at any site (Carrasco & Sarazin 2003; Sarazin & Tokovinin 2001) the mistaken idea of identifying high altitude winds with turbulence has become increasingly popular among the astronomical community. If there is indeed a relationship between V_0 and V_{200} at the different sites, this does not mean that the level of turbulence depends on V_{200} but only that the coherence time will vary with V_{200} behaviour. We wish to alert the astronomical community concerning this false interpretation of V_{200} as a parameter related to turbulence.

5 SUMMARY AND CONCLUSIONS

We have analysed the wind vertical profiles for the period 1980–2002 using the NCEP/NCAR Reanalysis archive at five astronomical sites. We have verified the consistency of the statistical results derived from these databases and balloon measurements. Our main results and conclusions may be summarized as follows:

1. The excellent statistical correlation between NCEP/NCAR Reanalysis data and balloon measurements reveal the NCEP/NCAR Reanalysis archive to be a useful source of meteorological data for site characterization.
2. In terms of high altitude wind speed (V_{200}) statistics, La Palma is the best of the five site studied, with a mean value of 22.13 m s^{-1} , the lowest standard deviation and the most stable V_{200} behaviour. La Silla is at the bottom of the ranking with a mean V_{200} of 33.35 m s^{-1} , although the seasonal V_{200} amplitude is the lowest of the five cases, with a statistical value of 12.46 m s^{-1} .
3. We have found a clear annual periodicity of V_{200} at the five observatories.
4. We have found a good level of correlation between high and low altitude winds, supporting a linear relationship between V_{200} and the average velocity of the turbulence at any of the sites. However, the proportional coefficient of this relationship may differ from site to site as a result of the clear seasonal behaviour of winds and the large differences of wind vertical profiles behaviour at the sites.

The results in this paper does not confirm a relationship of V_0 and V_{200} at the studied sites, but only a connection of high and low altitude winds. V_0 strongly depends on turbulence structure, and therefore, its connection to high altitude winds could be only establish with a proper turbulence characterisation. We alert the astronomical community on the wrong idea of identify high altitude winds and turbulence at the ground level that it is being more and more popular in the recent meetings. If a V_0 and V_{200} relationship were valid elsewhere, we may connect coherence time and high altitude winds, but never image quality.

ACKNOWLEDGMENTS

The authors thank T. Mahoney for his useful discussions and assistance. This work has made use of the NCEP Reanalysis data provided by the NOAA-CIRES Climate Diagnos-

tics Center, Boulder, Colorado, USA, from their Web site at <http://www.cdc.noaa.gov/>. Wavelet software was provided by C. Torrence and G. Compo, and is available at URL: <http://paos.colorado.edu/research/wavelets/>. This work has been partially funded by the Spanish Ministerio de Ciencia y Tecnologia (AYA2003-07728).

REFERENCES

- Carrasco, E. & Sarazin, M. 2003, RMxAA, 19, 103
- Chueca, S., García-Lorenzo, B., Muñoz-Tuñón, C., & Fuensalida, J.J. 2004, MNRAS, 349, 627
- García-Lorenzo, B., Fuensalida, J.J., & Eff-Darwich, A. 2004, SPIE Conf. on Remote Sensing, Maspalomas, Spain, SPIE, 5372, in press
- Kalnay, R., et al. 1996, The NCEP/NCAR 40-year Reanalysis Project, Bull. Am. Meteorol. Soc., 77, 437-471
- Kistler, R., et al. 2001, The NCEP-NCAR 50-Year reanalysis: Monthly means CD-ROM and documentation, Bull. Am. Meteorol. Soc., 82, 247-268, 2001
- Muñoz-Tuñón, C., Varela, A. & Mahoney, T. 1998, New Astron. Rev., 42, 409
- Sarazin, M., Graham, E., Beniston, M., & Riemann, M. 2003, SPIE Conf. on Astronomical Telescope and Instrumentation, Waikoloa, Hawaii, USA, SPIE, 4840, 54
- Sarazin, M., Beniston, M., Graham, E., & Riemann, M. 2002, SF2A-2002: Semaine de l'Astrophysique Francaise, 15
- Sarazin, M. 2002, ESPAS Site Summary Series: Mauna Kea, Issue 1.1
- Sarazin, M., & Tokovinin, A. 2001, Beyond conventional Adaptive Optics, Venice
- Seidel, D.J., Ross, R.J., Angell, J.K., & Reid, G.C. 2001, Journal of Geophysical Research (JGR), 106, 7857
- Torrence, C., & Compo, G.P. 1998, Bull. Am. Meteorol. Soc., 79, 61-78
- Varela, A.M., Muñoz-Tuñón, C. & Gurtubai, A.G. 2001, Highlights of Spanish astrophysics II, conf. 301, Edited by Jaime Zamorano, Javier Gorgas, and Jesus Gallego